

Prediction and forecast of daily PM10 concentrations in Brno and Graz

Ernst Stadlober

Graz University of Technology

Zuzana Hüberová, Jaroslav Michálek

Brno University of Technology

Österr. Statistiktage 2011, Graz

9. September 2011

Contents

Situation in Graz and Brno

- Meteorological and anthropogenic factors
- Measuring stations in Graz and Brno

Statistical Prediction Models

- Multiple Linear Regression
- Generalized Linear Model

Quality of the Models

The general Situation

Graz (263k inhabitants): Basin Area

- Low wind velocities, low precipitation, temperature inversion
- Traffic, domestic fuel

Brno (430k inhabitants): Open Basin Area

- Concentration of industry, motor traffic, heating plants
- Temperature inversion

Consequence


- Threshold value daily mean $PM_{10} \geq 50 \mu g/m^3$ exceeded regularly
- Graz: 36 – 105 exceedances in winter seasons
- Brno: 55 – 70 exceedances

Stations in Graz

Forecast: **Graz-Mitte** (traffic area), **Graz-Süd** (industrial zone)

Precipitation: Graz-Nord

Temperature inversion: Kalkleiten



Graz-Mitte: Inversion and Weekend Effect




Figure: Graz-Mitte: Five winter seasons 02/03–06/07


No Inversion: 40% lower than Inversion

Sun/Holiday: 30% lower than Working Day

Stations in Brno

Forecast: **Arboretum** (botanical garden)

Židenice (heavy traffic street), **Zvonařka** (traffic spot)



Training Data and Test Data for Graz

Model fits on Training Data

- **Graz-Mitte:** 7 winter seasons (23.10.02–31.3.09)
- **Graz-Süd:** 6 winter seasons (1.10.03–31.3.09)
- All models are fitted on **observed covariates**

Predictions (one-day forecasts) on Test Data

- **Graz-Mitte, Graz-Süd:** 1 winter season (3.11.09 - 31.3.10)

Covariates for Graz

- Ap_t : daily mean PM10 **day t**
- ΔT_t : mean temp diff to test point Kalkleiten **day t**
- V_t : mean wind speed **day t** (Graz Mitte for Graz Süd)
- $Prec_t$: 1 = prec., 0 = no prec. **on day t** (Graz Nord)
- Ap_{lag} : mean of PM10 (12.30 ($t - 2$) to 12.00 ($t - 1$))
- T_{lag} : categorized mean (temp means: 12.30 ($t - 2$) to 12.00 ($t - 1$)) (= 0 temp > 0, = 1 temp ≤ 0)
(Graz Mitte for Graz Süd)
- **saturday**: (0/1)
- **sunday**: (0/1)
- **febr**: (0/1)
- **march**: (0/1)

Training Data and Test Data for Brno

Model fits on Training Data

- Arboretum: 2 winter seasons 11/06–03/07, 10/07–03/08
- Židenice, Zvonarka: 1 winter season 11/07–03/08
- All models are fitted on predicted covariates

Predictions (one-day forecasts) on Test Data

- Arboretum, Židenice, Zvonarka:
1 winter season (1.10.08–31.3.09)

Covariates for Brno

- Ap_t : daily mean PM10 day t
- $V_t \sin D_t(\text{pred})$: mean of 4 predictions
(=velocity*sin(direction)) (0.00, 6.00, 12.00, 18.00 day t)
- $V_t \cos D_t(\text{pred})$: mean of 4 predictions
(=velocity*cos(direction))
- $T_t(\text{pred})$: mean of 4 predictions of temperature
- $Cover_t(\text{pred})$: mean of 4 predictions of coverage(values 1:10)
- Ap_{lag} : mean of PM10 (13.00 ($t - 2$) to 12.00 ($t - 1$))
- T_{lag} : mean of 4 measurements of temperature
(18.00 ($t - 2$), 0.00, 6.00, 12.00 ($t - 1$))

Covariates for Brno - 2

- HS_t : activity of heating plants day t
(Oct: 0.08, Nov: 0.14, Dec: 0.17,
Jan; 0.19, Feb: 0.16, Mar: 0.14)
- weekend: (0/1)
- sunday: (0/1)
- febr: (0/1)
- march: (0/1)

Multiple Linear Regression

Dependent variable $\sqrt{Ap_t}$

Model assumption

- $x_t^{(i)}$ ($i = 1, 2, \dots, m$) *metric input variables*
- $d_t^{(j)}$ ($j = 1, \dots, p$) *dummy (0/1) input variables at day t*
- *Linear model for $\sqrt{Ap_t}$:*

$$\sqrt{Ap_t} = \beta_0 + \sum_{i=1}^m \beta_i x_t^{(i)} + \sum_{j=1}^p \beta_{m+j} d_t^{(j)} + \epsilon_t \quad (1)$$

where $\epsilon_t \stackrel{iid}{\sim} N(0, \sigma^2)$.

Multiple Linear Regression: Graz-Mitte

| Variables x or d | beta | sd(beta) | t -value |
|----------------------|-------|----------|------------|
| const | 7.41 | 0.12 | 61.5 |
| ΔT_t | -0.30 | 0.018 | -16.6 |
| $A_{p_{lag}}$ | 0.02 | 0.001 | 15.1 |
| V_t | -1.59 | 0.132 | -12.0 |
| T_{lag} | 0.95 | 0.084 | 11.4 |
| sunday | -0.97 | 0.085 | -11.4 |
| march | 0.75 | 0.095 | 7.9 |
| saturday | -0.69 | 0.094 | -7.3 |
| $Prec_t$ | -0.51 | 0.071 | -7.1 |
| febr | 0.34 | 0.090 | 3.8 |
| | R^2 | se | n |
| | 0.63 | 1.10 | 1199 |

Multiple Linear Regression: Zvonařka in Brno

| Variables x or d | beta | sd(beta) | t-value |
|----------------------|-------|----------|---------|
| const | 8.78 | 0.543 | 16.2 |
| $V_t \sin D_t$ | 0.36 | 0.059 | 6.0 |
| Cover_t | -0.26 | 0.052 | -5.0 |
| march | -1.04 | 0.313 | -3.3 |
| T_{lag} | -0.19 | 0.058 | -3.2 |
| T_t | 0.17 | 0.062 | 2.7 |
| weekend | -0.63 | 0.242 | -2.6 |
| $A_{p\text{lag}}$ | 0.01 | 0.005 | 2.1 |
| $V_t \cos D_t$ | -0.08 | 0.056 | -1.5 |
| β_{45} | | 0.37 | |
| S | | 103 ° | |
| | R^2 | se | n |
| | 0.54 | 1.24 | 132 |

- HS_t , sunday, febr not significant

GLM model with gamma distribution and log link

Dependent variable Ap_t is gamma distributed

Model assumption

- $x_t^{(i)}$ ($i = 1, 2, \dots, m$) *metric input variables*
- $d_t^{(j)}$ ($j = 1, \dots, p$) *dummy (0/1) input variables*
- Ap_t has $\text{Gamma}(\phi\mu_t, 1/\phi)$ distribution, ϕ dispersion

$$\ln(\mu_t) = \ln(E(Ap_t|S_t)) = \beta_0 + \sum_{i=1}^m \beta_i x_t^{(i)} + \sum_{j=1}^p \beta_{m+j} d_t^{(j)}, \quad (2)$$

$E(Ap_t|S_t)$ cond. expect. of Ap_t for given set S_t of variables

GLM with gamma log link: Graz-Mitte

| Variables x or d | beta | sd(beta) |
|------------------------|--------|----------|
| const | 3.12 | 0.081 |
| ΔT_t | -0.08 | 0.005 |
| $\ln(Ap_{\text{lag}})$ | 0.31 | 0.020 |
| V_t | -0.50 | 0.038 |
| T_{lag} | 0.27 | 0.024 |
| sunday | -0.29 | 0.024 |
| march | 0.22 | 0.027 |
| saturday | -0.21 | 0.027 |
| $Prec_t$ | -0.15 | 0.021 |
| febr | 0.09 | 0.026 |
| χ^2_{ans} | 115.08 | |
| f | 1189 | |
| n | 1199 | |

GLM with gamma log link: Zvonařka in Brno

| Variables x or d | beta | sd(beta) |
|-----------------------|-------------|----------|
| const | 4.53 | 0.136 |
| $V_t \sin D_t$ | 0.12 | 0.017 |
| T_{lag} | -0.07 | 0.015 |
| Cover_t | -0.07 | 0.016 |
| march | -0.34 | 0.096 |
| T_t | 0.06 | 0.017 |
| weekend | -0.17 | 0.069 |
| febr | 0.18 | 0.090 |
| $V_t \cos D_t$ | -0.02 | 0.016 |
| $\hat{\beta}_S$ | 0.13 | |
| S | 101° | |
| χ^2_{ans} | 17.98 | |
| f | 125 | |
| n | 134 | |


- Ap_{lag} , HS_t , sunday not significant

Quality of forecasting

- **Training data set** to compute estimates $\hat{\beta}_i, \hat{\beta}_{j+m}$
- Model LM applied to **test data** yield estimates $\widehat{\sqrt{Ap_t}}$
- $\widehat{Ap_t} = (\widehat{\sqrt{Ap_t}})^2$
- Model GLM applied to **test data** yield estimates $\widehat{\ln(Ap_t)}$
- $\widehat{Ap_t} = \exp(\widehat{\ln(Ap_t)})$

Quality of forecasting - 2

Quality function $Q(x, y)$: (x = observation, y = forecast)



Quality: 5 levels green (1=excellent) to red (5=very bad)

Practical performance and comparison: Graz-Mitte




Figure: Graz-Mitte: Quality function of LM and GLM

Practical performance and comparison: Graz-Mitte - 2




Figure: Graz-Mitte: Forecasts $\widehat{Ap_t}$ of GLM vs LM

Practical performance and comparison: Graz Mitte - 3




Figure: Graz-Mitte: Quality function GLM vs LM

Practical performance and comparison: Graz Mitte - 5

Table: Contingency table Graz-Mitte: Quality of GLM (rows) vs LM (columns), no. and perc. of quality 1 to 5

| GLM/LM | 1 | 2 | 3 | 4 | 5 | sGLM | % | cum % |
|--------|------|------|------|------|-------|-------|------|-------|
| 1 | 63 | 5 | 0 | 0 | 0 | 68 | 45.6 | 45.6 |
| 2 | 4 | 27 | 5 | 0 | 1 | 37 | 24.8 | 70.5 |
| 3 | 1 | 3 | 16 | 2 | 0 | 22 | 14.8 | 85.2 |
| 4 | 0 | 0 | 3 | 10 | 1 | 14 | 9.4 | 94.6 |
| 5 | 0 | 0 | 0 | 1 | 7 | 8 | 5.4 | 100.0 |
| sLM | 68 | 35 | 24 | 13 | 9 | 149 | | |
| % | 45.6 | 23.5 | 16.1 | 8.7 | 6.0 | 100.0 | | |
| cum % | 45.6 | 69.1 | 85.2 | 94.0 | 100.0 | | | |

Practical performance and comparison: Zvonarka





Figure: Zvonarka: Quality function of LM and GLM

Practical performance and comparison: Zvonarka - 2




Figure: Zvonarka: Forecasts \widehat{Ap}_t of GLM vs LM

Practical performance and comparison: Zvonařka - 3




Figure: Zvonařka: Quality function GLM vs LM

Practical performance and comparison: Zvonařka - 4



Table: Contingency table Zvonařka: Quality of GLM (rows) vs LM (columns), no. and perc. of quality 1 to 5

| GLM/LM | 1 | 2 | 3 | 4 | 5 | sGLM | % | cum % |
|--------|------|------|------|------|-------|-------|------|-------|
| 1 | 61 | 11 | 0 | 0 | 0 | 72 | 40.7 | 40.7 |
| 2 | 7 | 20 | 6 | 1 | 0 | 34 | 19.2 | 59.9 |
| 3 | 2 | 3 | 12 | 1 | 2 | 20 | 11.3 | 71.2 |
| 4 | 1 | 1 | 2 | 14 | 2 | 20 | 11.3 | 82.5 |
| 5 | 0 | 2 | 1 | 9 | 19 | 31 | 17.5 | 100.0 |
| sLM | 71 | 37 | 21 | 25 | 23 | 177 | | |
| % | 40.1 | 20.9 | 11.9 | 14.1 | 13.0 | 100.0 | | |
| cum % | 40.1 | 61.0 | 72.9 | 87.0 | 100.0 | | | |

Practical Experience Graz-Mitte and Graz-Süd

- Daily predictions of PM10 since winter season 04/05
- Meteorological forecasts by ZAMG Styria till 11 in the morning
- PM10-Prediction during 1 and 3 PM at www.oekostadt.graz.at
- Daily update: comparison of predictions with measured values (day $t - 2$)
- e-mail with excel-file to Environmental Department of Graz
- 80 – 90% of the predictions at least satisfactory

Prediction Graz-Mitte, Graz-Süd 2010/2011



Conclusion

- LM and GLM models suitable for daily PM10 forecasts
- Results of models similar and comparable
- Brno: 1 or 2 seasons for training data, test data 2008/09
- Graz: 6 or 7 seasons for training data, test data 2009/10
- Brno: no measures for precipitation and inversion for each site specific selection of covariates
- Graz: all 9 covariates are significant
- Very Good quality: Arboretum (98% satisfactory)
- Good quality: Graz-Mitte, Graz-Süd (85%)
- Moderate quality: Zvonarka, Židenice (74%)